

SPOTLIGHT

Animal models for studying neural crest development: is the mouse different?

Elias H. Barriga¹, Paul A. Trainor^{2,3}, Marianne Bronner^{4,*} and Roberto Mayor^{1,*}

ABSTRACT

The neural crest is a uniquely vertebrate cell type and has been well studied in a number of model systems. Zebrafish, *Xenopus* and chick embryos largely show consistent requirements for specific genes in early steps of neural crest development. By contrast, knockouts of homologous genes in the mouse often do not exhibit comparable early neural crest phenotypes. In this Spotlight article, we discuss these species-specific differences, suggest possible explanations for the divergent phenotypes in mouse and urge the community to consider these issues and the need for further research in complementary systems.

KEY WORDS: Neural crest, Mouse, Zebrafish, *Xenopus*, Chicken, Gene knockout

Introduction

The neural crest is a multipotent stem cell population unique to vertebrates that contributes to development of the peripheral nervous system, craniofacial skeleton, adrenal gland, cardiac outflow tract, enteric ganglia of the gut, and other tissues (LeDouarin, 1982; Bronner and LeDouarin, 2012; Bronner, 2012; Mayor and Theveneau, 2013; Bhatt et al., 2013). The multipotency and migratory ability of the neural crest render it a cell type of great interest for regenerative medicine, as a therapeutic target for craniofacial defects (Trainor and Andrews, 2013) and for providing insights into vertebrate evolution (Simões-Costa and Bronner, 2013; Parker et al., 2014).

Many different vertebrates have emerged for studying neural crest biology in the past century: first amphibians and birds, followed by mouse, teleosts and, more recently, basal vertebrates such as lamprey and hagfish (Aybar and Mayor, 2002; Trainor, 2005; Sauka-Spengler et al., 2007; Ota et al., 2007). Loss-of-function analyses in animal models (primarily chick, *Xenopus* and zebrafish) have established a pan-vertebrate neural crest gene regulatory network (GRN), in which inhibition of individual transcriptional components leads to severe neural crest induction phenotypes (Simões-Costa and Bronner, 2015). Surprisingly, however, mouse mutants of the same genes often lack comparable phenotypes, at least at early stages of neural crest development. Here, we explore possible reasons for these apparent discrepancies between mouse and non-mammalian models.

Methodologies to study genes involved in neural crest development

Dominant-negative constructs, antisense morpholinos and RNAi have been the methods of choice for transiently perturbing neural crest development in chick, *Xenopus* and zebrafish embryos. In addition, several zebrafish neural crest mutants have emerged from forward genetic screens (Chakrabarti et al., 1983; Driever et al., 1996; Haffter et al., 1996). These loss-of-function approaches have yielded largely similar neural crest phenotypes and congruity, with only relatively minor differences, across non-mammalian species. Although TALENs and CRISPR-Cas9 are currently being employed across a wide variety of species to generate knockout animals (Peng et al., 2014), it is currently unknown whether they will generate similar results in neural crest studies across multiple species.

The mouse offers a powerful genetic model for studying gene function during development. Indeed, many mouse mutants exhibit neural crest phenotypes, particularly during craniofacial development, leading to important discoveries linked to human craniofacial malformations (Chai and Maxson, 2006; Sakai and Trainor, 2009; Trainor and Andrews, 2013; He and Soriano, 2013; Fantauzzo and Soriano, 2014). Strategies for generating mutants include spontaneous or radiation-induced mutations and gene targeting to produce null mice or conditional knockouts, using Cre/loxP or Flp/FRT technology.

There are several examples in which null mice completely lacking neural crest genes (e.g. *Fgf8*, *Pax3*, *Pax7*, *Notch1*) do not exhibit a failure of neural crest cell induction comparable to those observed in non-mammalian species (Conway et al., 1997; Frank et al., 2002). We describe potential reasons for these differences further below. To study mouse neural crest development in a spatiotemporal manner, the most widely used genetic tool for conditional gene knockout has been Wnt1-Cre (Danielian et al., 1998). However, this may not be ideal for early neural crest development for several reasons. First, Wnt1 expression is not restricted to the neural crest domain but rather marks dorsal neural stem cells that contribute to both central nervous system and neural crest progenitors (McMahon et al., 1992). Second, in avian embryos, expression of Wnt1 commences well after neural crest induction (Basch et al., 2006). If the same is true in mice (which has not yet been determined), Wnt1-Cre may initiate too late to elicit a neural crest induction phenotype. Third, the Wnt1-Cre line was recently found to elicit activation of Wnt signalling in ectopic locations (Lewis et al., 2013), particularly the midbrain, which might affect the interpretation of results related to the cranial neural crest. This is particularly important given the well-established role of Wnt signalling throughout neural crest development. Therefore, it will be important to develop and use other Cre driver lines to study neural crest induction in mice.

Differences in loss-of-function phenotypes between mouse and non-mammalian models

Clear discrepancies exist in the loss-of-function phenotypes observed between non-mammalian and mouse models.

¹Department of Cell and Developmental Biology, University College London, Gower Street, London WC1E 6BT, UK. ²Stowers Institute for Medical Research, Kansas City, MO 64110, USA. ³Department of Anatomy and Cell Biology, University of Kansas Medical Centre, Kansas City, KS 66160, USA. ⁴Division of Biology, California Institute of Technology, Pasadena, CA 91125, USA.

*Authors for correspondence (mbronner@caltech.edu; r.mayor@ucl.ac.uk)

Table 1. Comparison of early neural crest (NC) phenotypes generated in loss-of-function experiments between non-mammalian and mouse models

Gene/pathway	NC phenotype in non-mammalian models	NC phenotype in mouse
NC inductive factors		
Wnt (canonical)		
<i>Wnt1</i>	Impaired NC induction in <i>Xenopus</i> (mRNA [1]) and chick ($\Delta^{-/-}$ [2])	No NC induction phenotype (KO [3,4])
<i>Wnt3a</i>	Impaired NC induction in <i>Xenopus</i> (MO [5]) and impaired cardiac NC formation in zebrafish (MO, mRNA [6]); NC expansion in chick and zebrafish gain-of-function experiments (mRNA [6,7])	No NC induction phenotype (KO [4,8-13])
<i>Wnt8</i>	Impaired NC induction in <i>Xenopus</i> ($\Delta^{-/-}$ [14,15]) and zebrafish (MO [16])	No NC induction phenotype (KO [4])
<i>Wnt1/Wnt3a</i>	NA	No NC induction phenotype, late effect on NC maintenance (KO [3])
Retinoic acid (RARs)		
	Impaired NC induction in <i>Xenopus</i> ($\Delta^{-/-}$ [17]) and in quail ($\mu^{-/-}$ [18])	No NC induction phenotype, late effect on NC migration (triple KO [19])
<i>Fgf8</i>	Impaired NC induction in <i>Xenopus</i> ($\Delta^{-/-}$ [20]; MO [21]) and cranial patterning defects in zebrafish ($\mu^{-/-}$ [22])	No NC induction phenotype, late effect on migratory cardiac NC maintenance (KO [23])
Notch	Impaired NC induction in <i>Xenopus</i> ($\Delta^{-/-}$ [24]), chick ($\Delta^{-/-}$ [25]) and zebrafish ($\mu^{-/-}$ [26,27])	No NC induction phenotype, late effect on NC migration, proliferation and differentiation (KO ^{wc} [28])
NC GRN		
SoxE		
<i>Sox8</i>	Impaired NC induction in <i>Xenopus</i> (MO [29]) and chick (MO [30])	No NC induction phenotype (KO [31])
<i>Sox9</i>	Impaired NC induction in zebrafish (MO+ $\mu^{-/-}$ [32]) and <i>Xenopus</i> (MO [33,34]); impaired NC maintenance in chick (mRNA [35]; $\Delta^{-/-}$ [36]; MO [37])	No NC induction phenotype, late effect on migration and differentiation (KO ^{wc} [38,39])
<i>Tfap2</i>	Impaired NC induction in zebrafish ($\mu^{-/-}$ [40]), <i>Xenopus</i> (MO [41,42])	No NC induction phenotype, late effect on NC differentiation (KO [43,44])
<i>Foxd3</i>	Impaired NC induction in zebrafish (MO [45]; $\mu^{-/-}$ [46]), <i>Xenopus</i> ($\Delta^{-/-}$ [47,48]) and chick (MO [49])	No NC induction phenotype, late effect on NC maintenance (KO ^{wc} [50])
<i>Pax3/7</i>	Impaired NC induction in <i>Xenopus</i> (MO [51]) and chick (MO [52]); partial inhibition of NC induction in zebrafish (MO [53])	No NC induction phenotype with respect to cranial and cardiac NC, but impaired trunk NC induction; late effect on cardiac NC migration ($\mu^{-/-}$ [54])
<i>Msx1/2</i>	Impaired NC induction in <i>Xenopus</i> ($\Delta^{-/-}$ [55]); impaired NC induction in zebrafish when both <i>msxb</i> and <i>msxc</i> are inhibited (MO [56]); NC expansion in chick gain of function [57,58]	No NC induction phenotype, late effect on NC maintenance and differentiation in the double KO <i>Msx1^{-/-};Msx2^{-/-}</i> [59]
<i>Snail1/2</i>	Impaired NC induction in <i>Xenopus</i> ($\Delta^{-/-}$ [60,61]; MO [62]); early migration phenotype in chick (AS [63]; MO [64])	No NC induction phenotype, early lethality (KO [65,66])
NC migration factors		
Wnt (non-canonical)		
<i>Wnt11</i>	Impaired early NC migration in zebrafish ($\mu^{-/-}$ [67]) and <i>Xenopus</i> (MO [68,69])	No NC migration phenotype (KO [4,70])
<i>Dvl</i>	Impaired early NC migration in zebrafish ($\Delta^{-/-}$ [67]), <i>Xenopus</i> (MO [69]; $\Delta^{-/-}$ [71]) and chick ($\Delta^{-/-}$ [72])	No early NC migration phenotype, late effect on cardiac NC differentiation (KO [4,73])
<i>Vangl1/2</i>	Impaired NC migration in zebrafish ($\mu^{-/-}$, $\Delta^{-/-}$ [74]) and <i>Xenopus</i> (MO [69])	No NC induction or migration phenotype (KO, KO ^{wc} [75])
<i>Sdc4</i>	Impaired NC migration in zebrafish and <i>Xenopus</i> (MO [74])	No NC induction or migration phenotype, late effect on cranial cartilage (KO [76])
<i>N-cad/Cdh2</i>	Impaired NC migration in zebrafish (MO, $\mu^{-/-}$ [77]), <i>Xenopus</i> (MO [78]) and chick (BA [79]; $\Delta^{-/-}$ [80])	No early NC migration phenotype, late effect on cardiac NC differentiation (KO ^{wc} [81])

These are examples of genes that show a clear discrepancy in loss-of-function phenotypes between non-mammalian (in at least two species) and mouse. Many other examples that show a similar discrepancy with mouse are not included here because they have been studied in only one non-mammalian species. MO, morpholino; AS, antisense oligonucleotide; $\mu^{-/-}$ or $\mu^{+/-}$, homozygous or heterozygous mutant; $\Delta^{-/-}$, dominant negative; KO, full knockout; KO^{wc}, Wnt1-Cre KO; BA, blocking antibody; mRNA, gain-of-function. NA, not applicable. RAR, retinoic acid receptor.

References: [1] Saint-Jeannet et al., 1997; [2] Garcia-Castro et al., 2002; [3] Ikeya et al., 1997; [4] van Amerongen and Berns, 2006; [5] Elkouby et al., 2010; [6] Sun et al., 2008; [7] Patthey et al., 2009; [8] Takada et al., 1994; [9] Greco et al., 1996; [10] Yoshikawa et al., 1997; [11] Lee et al., 2000; [12] Ikeya and Takada, 2001; [13] Nakaya et al., 2005; [14] LaBonne and Bronner-Fraser, 1998; [15] Steventon et al., 2009; [16] Lewis et al., 2004; [17] Villanueva et al., 2002; [18] Martínez-Morales et al., 2011; [19] Dupé and Pellerin, 2009; [20] Mayor et al., 1997; [21] Monsoro-Burq et al., 2005; [22] Roehl and Nüsslein-Volhard, 2001; [23] Frank et al., 2002; [24] Glavic et al., 2004; [25] Endo et al., 2002; [26] Jiang et al., 1996; [27] Cornell and Eisen, 2002; [28] Mead and Yutzey, 2012; [29] O'Donnel et al., 2006; [30] Betancur et al., 2011; [31] Sock et al., 2001; [32] Yan et al., 2005; [33] Aoki et al., 2003; [34] Spokony et al., 2002; [35] Cheung and Briscoe, 2003; [36] Cheung et al., 2005; [37] Betancur et al., 2010b; [38] Mori-Akiyama et al., 2003; [39] Akiyama et al., 2004; [40] Knight et al., 2003; [41] Luo et al., 2003; [42] Hong et al., 2014; [43] Schorle et al., 1996; [44] Zhang et al., 1996; [45] Lister et al., 2006; [46] Stewart et al., 2006; [47] Pohl and Knochel, 2001; [48] Sasai et al., 2001; [49] Fairchild et al., 2014; [50] Teng et al., 2008; [51] Maczkowiak et al., 2010; [52] Basch et al., 2006; [53] Minchin and Hughes, 2008; [54] Conway et al., 1997; [55] Tribulo et al., 2003; [56] Phillips et al., 2006; [57] Liu et al., 2004; [58] Barembaum and Bronner, 2013; [59] Ishii et al., 2005; [60] LaBonne and Bronner-Fraser, 2000; [61] Aybar et al., 2002; [62] Shi et al., 2011; [63] Nieto et al., 1994; [64] Taneyhill et al., 2007; [65] Jiang et al., 1998; [66] Murray and Gridley, 2006; [67] Banerjee et al., 2011; [68] Matthews et al., 2008a; [69] Carmona-Fontaine et al., 2008; [70] Majumdar et al., 2003; [71] De Calisto et al., 2005; [72] Rios et al., 2011; [73] Hamblet et al., 2002; [74] Matthews et al., 2008b; [75] Pryor et al., 2014; [76] Echtermeyer et al., 2001; [77] Piloto and Schilling, 2010; [78] Theveneau et al., 2010; [79] Bronner-Fraser et al., 1992; [80] Nakagawa and Takeichi, 1998; [81] Luo et al., 2006.

Summarised in Table 1 are those cases in which the same gene has been studied in more than one non-mammalian species. Key examples are discussed below.

There is strong evidence that canonical Wnt signalling is essential for neural crest induction. Although the requirement of a specific Wnt ligand can vary among species, the inhibition of Wnt signalling, their receptors or downstream components has a dramatic effect on neural crest formation in chick, *Xenopus* and zebrafish (Saint-Jeannet et al., 1997; LaBonne and Bronner-Fraser, 1998; Garcia-Castro et al., 2002; Lewis et al., 2004; Sun et al., 2008; Patthey et al., 2009; Steventon et al., 2009; Elkouby et al., 2010). Surprisingly, however, knockout of canonical Wnt genes in mouse does not affect neural crest induction (Takada et al., 1994; Greco et al., 1996; Yoshikawa et al., 1997; Ikeya et al., 1997; Lee et al., 2000; Ikeya and Takada, 2001; Rodríguez-Mari et al., 2005; Nakaya et al., 2005; van Amerongen and Berns, 2006).

During the induction process, the neural crest GRN becomes activated (Betancur et al., 2010a; Steventon et al., 2005). This transcriptional cascade is highly conserved, even in the basal jawless vertebrate lamprey (Sauka-Spengler et al., 2007; Nikitina and Bronner-Fraser, 2009; Nikitina et al., 2008). Two of the key genes are *Snail1/2* and *Foxd3*, both of which are required for neural crest induction in zebrafish, *Xenopus* and chick (Nieto et al., 1994; LaBonne and Bronner-Fraser, 2000; Pohl and Knöchel, 2001; Sasai et al., 2001; Aybar et al., 2003; Lister et al., 2006; Stewart et al., 2006; Taneyhill et al., 2007; Shi et al., 2011; Fairchild et al., 2014). However, neural crest induction is unaffected in *Snail1/2* or *Foxd3* knockout mice (Jiang et al., 1998; Murray and Gridley, 2006; Teng et al., 2008), although *Foxd3* knockouts exhibit a defect in late neural crest maintenance (Teng et al., 2008).

A defining feature of the neural crest is its migratory capacity. Many of the cellular mechanisms underlying migration are conserved across species; for example, in *Xenopus*, zebrafish and chick, non-canonical Wnt signalling is essential for normal neural crest migration (De Calisto et al., 2005; Carmona-Fontaine et al., 2008; Matthews et al., 2008a; Matthews et al., 2008b; Rios et al., 2011; Banerjee et al., 2011). By contrast, inhibition of non-canonical Wnt signalling in mouse does not affect migration (Majumdar et al., 2003; van Amerongen and Berns, 2006; Pryor et al., 2014).

Thus, key factors implicated in the induction and migration steps of neural crest development in non-mammalian species appear to be dispensable for those stages in mouse, at least as assayed by the most commonly used methodologies. This raises the intriguing possibility that the mechanisms of early neural crest development are divergent in the mouse lineage, or that there is a higher redundancy of genes in mouse compared with non-mammalian species. These and other alternatives are discussed below.

Possible explanations for the differences between mouse and other vertebrates

As detailed in Table 1, genes required for different steps of early neural crest development, from induction to migration, are largely conserved between avians, amphibians and fish, whereas specific knockouts of homologous mouse genes only have later neural crest defects. Several possible explanations may account for these differences.

First, there may be limited conservation of neural crest genes between mouse and non-mammalian organisms. It is well known that there is a greater temporal separation between neurulation and neural crest cell induction in mice than in frogs, fish and (to a lesser

extent) avians. Thus, there might be fundamental differences in the temporal induction of neural crest cells in non-mammalian versus mammalian species, perhaps accompanied by differences in the neural crest GRN. However, given that the expression and function of neural crest GRN components are conserved between fish, frog, birds and lamprey (Simões-Costa and Bronner, 2013), this seems unlikely and non-parsimonious. The relevance of these differences, if any, should be further explored to gain a deeper understanding of the conservation and diversity of neural crest cell induction mechanisms across species. This highlights the importance of developing other mammalian model systems, such as the rabbit or pig (Vadasz et al., 2013).

Second, the Wnt1-Cre line might be inappropriate to study early stages of neural crest development, as the initiation of neural crest development, at least in non-mammalian species, occurs prior to the initiation of Wnt1 expression. In addition, even though Wnt1-Cre lineage tracing labels most neural crest cells in mice, if a gene of interest or prospective neural crest cell is present in a domain that is broader than that demarcated by Wnt1, then Wnt1-Cre may not excise the gene completely and adjacent cells might compensate. Thus, it is crucial to explore these issues further during mouse development to ascertain the degree of conservation and diversity of neural crest induction across species.

Third, technical limitations of the Wnt1-Cre line, such as ectopic activation of Wnt signalling, may confound some analyses, as increasing Wnt signalling could promote neural crest induction and thereby rescue the phenotype.

Fourth, there might be redundancy or compensation in the mouse. Many neural crest GRN transcription factors have been duplicated during evolution, perhaps leading to paralogous factors or the co-option of members of other gene families assuming the same function in the GRN. Like all extant models, the mouse is derived and has a particularly long phase of early development, perhaps enabling additional gene compensation and/or rewiring of its neural crest GRN.

Finally, non-specific off-target effects from transient knockdown techniques may affect the interpretation of phenotypes. Use of morpholinos or dominant negatives is less specific than the production of null or conditional mice, perhaps partially accounting for differences in phenotypes. However, the reproducibility of most phenotypes across species, together with important controls such as rescue experiments and comparison with zebrafish mutants where they exist, makes this less likely.

Conclusions

Regardless of the underlying reasons for the differences between mouse and other species, we recommend exercising caution in interpreting negative results from knockout mice regarding a role in early neural crest development. The absence of an obvious phenotype does not necessarily mean that the gene of interest is not normally involved in the process. It could highlight a redundant function in mice rather than a fundamental difference with other species. To date, it is unknown whether differences between mouse and non-mammalian species reflect peculiarities of mouse development or are generally true for mammals. This is a particularly important question because the mouse is typically considered a better model for human development (and developmental disease) than non-mammalian species. Future neural crest studies using other mammalian species are therefore needed to clarify the degree of conservation within the mammalian lineage, and hopefully to provide insight into human neural crest development and defects thereof.

In the mouse, studies using a new Wnt1-Cre line that corrects ectopic expression (Lewis et al., 2013) will be very informative, as will the development of other Cre drivers that function earlier in development at epiblast stages and are restricted to the prospective neural crest; hence, considerable attention from the field needs to be devoted to developing new spatiotemporally appropriate lines. Equally, transient knockdown techniques suffer from problems including dilution over time and a requirement for careful controls to guarantee specificity. In the long term, analogous knockdown experiments must be performed across numerous species, including other mammals, to systematically examine the role of a gene of interest and its paralogues. The recent advent of CRISPR-Cas9 will facilitate comparative studies of this kind across a range of important vertebrates to definitively establish key players in neural crest development and GRN changes during vertebrate evolution. This will help establish the degree of redundancy versus species specificity in the molecular and cellular mechanisms governing neural crest induction and migration. Although an overlap of core GRNs components is expected, there might also be many surprises.

The availability of new technologies that enable analysis across a wide spectrum of species makes for exciting times in the field. Such studies are essential to determine which animal model or combination of models is best suited for understanding neural crest development, evolution and disease.

Competing interests

The authors declare no competing or financial interests.

Funding

R.M. is supported by grants from the Medical Research Council [MR/J000655/1] and Wellcome Trust. E.H.B. is supported by a Quantitative Fellowship [4505]. M.B. is supported by National Institutes of Health grants [DE024157 and HD037105]. P.A.T. is supported by the Stowers Institute for Medical Research and the National Institute of Dental and Craniofacial Research [DE 016082].

References

- Akiyama, H., Chaboissier, M.-C., Behringer, R. R., Rowitch, D. H., Schedl, A., Epstein, J. A. and de Crombrughe, B. (2004). Essential role of Sox9 in the pathway that controls formation of cardiac valves and septa. *Proc. Natl. Acad. Sci. USA* **101**, 6502–6507.
- Aoki, Y., Saint-Germain, N., Gyda, M., Magner-Fink, E. K., Lee, Y.-H., Credidio, C. and Saint-Jeannet, J.-P. (2003). Sox10 regulates the development of neural crest-derived melanocytes in *Xenopus*. *Dev. Biol.* **259**, 19–33.
- Aybar, M. J. and Mayor, R. (2002). Early induction of neural crest cells: lessons learned from frog, fish and chick. *Curr. Opin. Genet. Dev.* **12**, 452–458.
- Aybar, M. J., Nieto, M. A. and Mayor, R. (2003). Snail precedes slug in the genetic cascade required for the specification and migration of the *Xenopus* neural crest. *Development* **130**, 483–494.
- Banerjee, S., Gordon, L., Donn, T. M., Berti, C., Moens, C. B., Burden, S. J. and Granato, M. (2011). A novel role for MuSK and non-canonical Wnt signaling during segmental neural crest cell migration. *Development* **138**, 3287–3296.
- Barenbaum, M. and Bronner, M. E. (2013). Identification and dissection of a key enhancer mediating cranial neural crest specific expression of transcription factor, *Ets-1*. *Dev. Biol.* **382**, 567–575.
- Basch, M. L., Bronner-Fraser, M. and García-Castro, M. I. (2006). Specification of the neural crest occurs during gastrulation and requires Pax7. *Nature* **441**, 218–222.
- Betancur, P., Bronner-Fraser, M. and Sauka-Spengler, T. (2010a). Assembling neural crest regulatory circuits into a gene regulatory network. *Annu. Rev. Cell Dev. Biol.* **26**, 581–603.
- Betancur, P., Bronner-Fraser, M. and Sauka-Spengler, T. (2010b). Genomic code for Sox10 activation reveals a key regulatory enhancer for cranial neural crest. *Proc. Natl. Acad. Sci. USA* **107**, 3570–3575.
- Betancur, P., Sauka-Spengler, T. and Bronner, M. (2011). A Sox10 enhancer element common to the otic placode and neural crest is activated by tissue-specific paralogs. *Development* **138**, 3689–3698.
- Bhatt, S., Diaz, R. and Trainor, P. A. (2013). Signals and switches in mammalian neural crest cell differentiation. *Cold Spring Harb. Perspect. Biol.* **5**, a008326.
- Bronner, M. E. (2012). Formation and migration of neural crest cells in the vertebrate embryo. *Histochem. Cell Biol.* **138**, 179–186.
- Bronner, M. E. and LeDouarin, N. M. (2012). Development and evolution of the neural crest: an overview. *Dev. Biol.* **366**, 2–9.
- Bronner-Fraser, M., Wolf, J. J. and Murray, B. A. (1992). Effects of antibodies against N-cadherin and N-CAM on the cranial neural crest and neural tube. *Dev. Biol.* **153**, 291–301.
- Carmona-Fontaine, C., Matthews, H. K., Kuriyama, S., Moreno, M., Dunn, G. A., Parsons, M., Stern, C. D. and Mayor, R. (2008). Contact inhibition of locomotion in vivo controls neural crest directional migration. *Nature* **456**, 957–961.
- Chai, Y. and Maxson, R. E., Jr (2006). Recent advances in craniofacial morphogenesis. *Dev. Dyn.* **235**, 2353–2375.
- Chakrabarti, S., Streisinger, G., Singer, F. and Walker, C. (1983). Frequency of gamma-ray induced specific locus and recessive lethal mutations in mature germ cells of the zebrafish, *Brachydanio rerio*. *Genetics* **103**, 109–123.
- Cheung, M. and Briscoe, J. (2003). Neural crest development is regulated by the transcription factor Sox9. *Development* **130**, 5681–5693.
- Cheung, M., Chaboissier, M.-C., Mynett, A., Hirst, E., Schedl, A. and Briscoe, J. (2005). The transcriptional control of trunk neural crest induction, survival, and delamination. *Dev. Cell* **8**, 179–192.
- Conway, S. J., Henderson, D. J. and Copp, A. J. (1997). Pax3 is required for cardiac neural crest migration in the mouse: evidence from the *spot* (*Sp^{2H}*) mutant. *Development* **124**, 505–514.
- Cornell, R. A. and Eisen, J. S. (2002). Delta/Notch signaling promotes formation of zebrafish neural crest by repressing Neurogenin 1 function. *Development* **129**, 2639–2648.
- Danielian, P. S., Muccino, D., Rowitch, D. H., Michael, S. K. and McMahon, A. P. (1998). Modification of gene activity in mouse embryos in utero by a tamoxifen-inducible form of Cre recombinase. *Curr. Biol.* **8**, 1323–1326.
- De Calisto, J., Araya, C., Marchant, L., Riaz, C. F. and Mayor, R. (2005). Essential role of non-canonical Wnt signalling in neural crest migration. *Development* **132**, 2587–2597.
- Driever, W., Solnica-Krezel, L., Schier, A. F., Neuhauss, S. C. F., Malicki, J., Stemple, D. L., Stainier, D. Y. R., Zwartkruis, F., Abdelilah, S., Rangini, Z. et al. (1996). A genetic screen for mutations affecting embryogenesis in zebrafish. *Development* **123**, 37–46.
- Dupé, V. and Pellerin, I. (2009). Retinoic acid receptors exhibit cell-autonomous functions in cranial neural crest cells. *Dev. Dyn.* **238**, 2701–2711.
- Echtermeyer, F., Streit, M., Wilcox-Adelman, S., Saoncella, S., Denhez, F., Detmar, M. and Goetinck, P. F. (2001). Delayed wound repair and impaired angiogenesis in mice lacking syndecan-4. *J. Clin. Invest.* **107**, R9–R14.
- Elkouby, Y. M., Elias, S., Casey, E. S., Blythe, S. A., Tsabar, N., Klein, P. S., Root, H., Liu, K. J. and Frank, D. (2010). Mesodermal Wnt signaling organizes the neural plate via Meis3. *Development* **137**, 1531–1541.
- Endo, Y., Osumi, N. and Wakamatsu, Y. (2002). Bimodal functions of Notch-mediated signaling are involved in neural crest formation during avian ectoderm development. *Development* **129**, 863–873.
- Fairchild, C. L., Conway, J. P., Schiffmacher, A. T., Taneyhill, L. A. and Gammill, L. S. (2014). FoxD3 regulates cranial neural crest EMT via downregulation of tetraspanin18 independent of its functions during neural crest formation. *Mech. Dev.* **132**, 1–12.
- Fantauzzo, K. A. and Soriano, P. (2014). PI3K-mediated PDGFR α signaling regulates survival and proliferation in skeletal development through p53-dependent intracellular pathways. *Genes Dev.* **28**, 1005–1017.
- Frank, D. U., Fotheringham, L. K., Brewer, J. A., Muglia, L. J., Tristani-Firouzi, M., Capecchi, M. R. and Moon, A. M. (2002). An Fgf8 mouse mutant phenocopies human 22q11 deletion syndrome. *Development* **129**, 4591–4603.
- García-Castro, M., Marcelle, C. and Bronner-Fraser, M. (2002). Ectodermal Wnt function as a neural crest inducer. *Science* **297**, 848–851.
- Glavic, A., Silva, F., Aybar, M. J., Bastidas, F. and Mayor, R. (2004). Interplay between Notch signaling and the homeoprotein Xir1 is required for neural crest induction in *Xenopus* embryos. *Development* **131**, 347–359.
- Greco, T. L., Takada, S., Newhouse, M. M., McMahon, J. A., McMahon, A. P. and Camper, S. A. (1996). Analysis of the vestigial tail mutation demonstrates that Wnt-3a gene dosage regulates mouse axial development. *Genes Dev.* **10**, 313–324.
- Haffter, P., Granato, M., Brand, M., Mullins, M. C., Hammerschmidt, M., Kane, D. A., Odenthal, J., van Eeden, F. J., Jiang, Y. J., Heisenberg, C. P. et al. (1996). The identification of genes with unique and essential functions in the development of the zebrafish, *Danio rerio*. *Development* **123**, 1–3.
- Hamblet, N. S., Lijam, N., Ruiz-Lozano, P., Wang, J., Yang, Y., Luo, Z., Mei, L., Chien, K. R., Sussman, D. J. and Wynshaw-Boris, A. (2002). Dishevelled 2 is essential for cardiac outflow tract development, somite segmentation and neural tube closure. *Development* **129**, 5827–5838.
- He, F. and Soriano, P. (2013). A critical role for PDGFR α signaling in medial nasal process development. *PLoS Genet.* **9**, e1003851.
- Hong, C.-S., Devotta, A., Lee, Y.-H., Park, B.-Y. and Saint-Jeannet, J.-P. (2014). Transcription factor AP2 epsilon (Tap2e) regulates neural crest specification in *xenopus*. *Dev. Neurobiol.* **74**, 894–906.
- Ikeya, M. and Takada, S. (2001). Wnt-3a is required for somite specification along the anteroposterior axis of the mouse embryo and for regulation of cdx-1 expression. *Mech. Dev.* **103**, 27–33.
- Ikeya, M., Lee, S. M. K., Johnson, J. E., McMahon, A. P. and Takada, S. (1997). Wnt signalling required for expansion of neural crest and CNS progenitors. *Nature* **389**, 966–970.

- Ishii, M., Han, J., Yen, H.-Y., Sucov, H. M., Chai, Y. and Maxson, R. E., Jr (2005). Combined deficiencies of *Msx1* and *Msx2* cause impaired patterning and survival of the cranial neural crest. *Development* **132**, 4937–4950.
- Jiang, Y., Brand, M., Heisenberg, C. P., Beuchle, D., Furutani-Seiki, M., Kelsh, R. N., Warga, R. M., Granato, M., Haffter, P., Hammerschmidt, M. et al. (1996). Mutations affecting neurogenesis and brain morphology in the zebrafish, *Danio rerio*. *Development* **123**, 205–216.
- Jiang, R., Lan, Y., Norton, C. R., Sundberg, J. P. and Gridley, T. (1998). The *Slug* gene is not essential for mesoderm or neural crest development in mice. *Dev. Biol.* **198**, 277–285.
- Knight, R. D., Nair, S., Nelson, S. S., Afshar, A., Javidan, Y., Geisler, R., Rauch, G.-J. and Schilling, T. F. (2003). *lockjaw* encodes a zebrafish *tfap2a* required for early neural crest development. *Development* **130**, 5755–5768.
- LaBonne, C. and Bronner-Fraser, M. (1998). Neural crest induction in *Xenopus*: evidence for a two-signal model. *Development* **125**, 2403–2414.
- LaBonne, C. and Bronner-Fraser, M. (2000). Snail-related transcriptional repressors are required in *Xenopus* for both the induction of the neural crest and its subsequent migration. *Dev. Biol.* **221**, 195–205.
- LeDouarin, N. (1982). *The Neural Crest*. New York, NY: Cambridge University Press.
- Lee, S. M. K., Tole, S., Grove, E. and McMahon, A. P. (2000). A local *Wnt-3a* signal is required for development of the mammalian Hippocampus. *Development* **127**, 457–467.
- Lewis, J. L., Bonner, J., Modrell, M., Ragland, J. W., Moon, R. T., Dorsky, R. I. and Raible, D. W. (2004). Reiterated *Wnt* signaling during zebrafish neural crest development. *Development* **131**, 1299–1308.
- Lewis, A. E., Vasudevan, H. N., O'Neill, A. K., Soriano, P. and Bush, J. O. (2013). The widely used *Wnt1-Cre* transgene causes developmental phenotypes by ectopic activation of *Wnt* signaling. *Dev. Biol.* **379**, 229–234.
- Lister, J. A., Cooper, C., Nguyen, K., Modrell, M., Grant, K. and Raible, D. W. (2006). Zebrafish *Foxd3* is required for development of a subset of neural crest derivatives. *Dev. Biol.* **290**, 92–104.
- Liu, Y., Helms, A. W. and Johnson, J. E. (2004). Distinct activities of *Msx1* and *Msx3* in dorsal neural tube development. *Development* **131**, 1017–1028.
- Luo, T., Lee, Y.-H., Saint-Jeannet, J.-P. and Sargent, T. D. (2003). Induction of neural crest in *Xenopus* by transcription factor *AP2alpha*. *Proc. Natl. Acad. Sci. USA* **100**, 532–537.
- Luo, Y., High, F. A., Epstein, J. A. and Radice, G. L. (2006). N-cadherin is required for neural crest remodeling of the cardiac outflow tract. *Dev. Biol.* **299**, 517–528.
- Mackowiak, F., Matéos, S., Wang, E., Roche, D., Harland, R. and Monsoro-Burq, A. H. (2010). The *Pax3* and *Pax7* paralogs cooperate in neural and neural crest patterning using distinct molecular mechanisms, in *Xenopus laevis* embryos. *Dev. Biol.* **340**, 381–396.
- Majumdar, A., Vainio, S., Kispert, A., McMahon, J. and McMahon, A. P. (2003). *Wnt11* and *Ret/Gdnf* pathways cooperate in regulating ureteric branching during metanephric kidney development. *Development* **130**, 3175–3185.
- Martínez-Morales, P. L., Díez del Corral, R., Olivera-Martínez, I., Quiroga, A. C., Das, R. M., Barbas, J. A., Storey, K. G. and Morales, A. V. (2011). FGF and retinoic acid activity gradients control the timing of neural crest cell emigration in the trunk. *J. Cell Biol.* **194**, 489–503.
- Matthews, H. K., Broders-Bondon, F., Thiery, J. P. and Mayor, R. (2008a). *Wnt11r* is required for cranial neural crest migration. *Dev. Dyn.* **237**, 3404–3409.
- Matthews, H. K., Marchant, L., Carmona-Fontaine, C., Kuriyama, S., Larrain, J., Holt, M. R., Parsons, M. and Mayor, R. (2008b). Directional migration of neural crest cells in vivo is regulated by *Syndecan-4/Rac1* and non-canonical *Wnt* signaling/*RhoA*. *Development* **135**, 1771–1780.
- Mayor, R. and Theveneau, E. (2013). The neural crest. *Development* **140**, 2247–2251.
- Mayor, R., Guerrero, N. and Martínez, C. (1997). Role of FGF and noggin in neural crest induction. *Dev. Biol.* **189**, 1–12.
- McMahon, A. P., Joyner, A. L., Bradley, A. and McMahon, J. A. (1992). The midbrain-hindbrain phenotype of *Wnt-1/Wnt-1-* mice results from stepwise deletion of engrailed-expressing cells by 9.5 days postcoitum. *Cell* **69**, 581–595.
- Mead, T. J. and Yutzy, K. E. (2012). Notch pathway regulation of neural crest cell development in vivo. *Dev. Dyn.* **241**, 376–389.
- Minchin, J. E. N. and Hughes, S. M. (2008). Sequential actions of *Pax3* and *Pax7* drive xanthophore development in zebrafish neural crest. *Dev. Biol.* **317**, 508–522.
- Monsoro-Burq, A. H., Wang, E. and Harland, R. (2005). *Msx1* and *Pax3* cooperate to mediate FGF8 and WNT signals during *Xenopus* neural crest induction. *Dev. Cell* **8**, 167–178.
- Mori-Akiyama, Y., Akiyama, H., Rowitch, D. H. and de Crombrughe, B. (2003). *Sox9* is required for determination of the chondrogenic cell lineage in the cranial neural crest. *Proc. Natl. Acad. Sci. USA* **100**, 9360–9365.
- Murray, S. A. and Gridley, T. (2006). Snail family genes are required for left-right asymmetry determination, but not neural crest formation, in mice. *Proc. Natl. Acad. Sci. USA* **103**, 10300–10304.
- Nakagawa, S. and Takeichi, M. (1998). Neural crest emigration from the neural tube depends on regulated cadherin expression. *Development* **125**, 2963–2971.
- Nakaya, M.-a., Biris, K., Tsukiyama, T., Jaime, S., Rawls, J. A. and Yamaguchi, T. P. (2005). *Wnt3a* links left-right determination with segmentation and anteroposterior axis elongation. *Development* **132**, 5425–5436.
- Nieto, M. A., Sargent, M. G., Wilkinson, D. G. and Cooke, J. (1994). Control of cell behavior during vertebrate development by *Slug*, a zinc finger gene. *Science* **264**, 835–839.
- Nikitina, N. and Bronner-Fraser, M. (2009). Gene regulatory networks that control the specification of neural-crest cells in the lamprey. *Biochim. Biophys. Acta* **1789**, 274–278.
- Nikitina, N., Sauka-Spengler, T., Bronner-Fraser, M. (2008). Dissecting early regulatory relationships in the lamprey neural crest gene network. *Proc. Natl. Acad. Sci. USA* **105**, 20083–20088.
- O'Donnell, M., Hong, C.-S., Huang, X., Delnicki, R. J. and Saint-Jeannet, J.-P. (2006). Functional analysis of *Sox8* during neural crest development in *Xenopus*. *Development* **133**, 3817–3826.
- Ota, K. G., Kuraku, S. and Kuratani, S. (2007). Hagfish embryology with reference to the evolution of the neural crest. *Nature* **446**, 672–675.
- Parker, H. J., Bronner, M. E. and Krumlauf, R. (2014). A *Hox* regulatory network of hindbrain segmentation is conserved to the base of vertebrates. *Nature* **514**, 490–493.
- Patthey, C., Edlund, T. and Gunhaga, L. (2009). *Wnt*-regulated temporal control of BMP exposure directs the choice between neural plate border and epidermal fate. *Development* **136**, 73–83.
- Peng, Y., Clark, K. J., Campbell, J. M., Panetta, M. R., Guo, Y. and Ekker, S. C. (2014). Making designer mutants in model organisms. *Development* **141**, 4042–4054.
- Phillips, B. T., Kwon, H.-J., Melton, C., Houghtaling, P., Fritz, A. and Riley, B. B. (2006). Zebrafish *msxB*, *msxC* and *msxE* function together to refine the neural–nonneural border and regulate cranial placodes and neural crest development. *Dev. Biol.* **294**, 376–390.
- Piloto, S. and Schilling, T. F. (2010). *Ovo1* links *Wnt* signaling with N-cadherin localization during neural crest migration. *Development* **137**, 1981–1990.
- Pohl, B. S. and Knöchel, W. (2001). Overexpression of the transcriptional repressor *FoxD3* prevents neural crest formation in *Xenopus* embryos. *Mech. Dev.* **103**, 93–106.
- Pryor, S. E., Massa, V., Savery, D., Andre, P., Yang, Y., Greene, N. D. E. and Copp, A. J. (2014). *Vangl*-dependent planar cell polarity signalling is not required for neural crest migration in mammals. *Development* **141**, 3153–3158.
- Rios, A. C., Serralbo, O., Salgado, D. and Marcelle, C. (2011). Neural crest regulates myogenesis through the transient activation of *NOTCH*. *Nature* **473**, 532–535.
- Rodríguez-Marí, A., Yan, Y.-L., BreMiller, R. A., Wilson, C., Cañestro, C. and Postlethwait, J. H. (2005). Characterization and expression pattern of zebrafish anti-Müllerian hormone (*amh*) relative to *sox9a*, *sox9b*, and *cyp19a1a*, during gonad development. *Gene Expr. Patterns* **5**, 655–667.
- Roehl, H. and Nüsslein-Volhard, C. (2001). Zebrafish *pea3* and *erm* are general targets of FGF8 signaling. *Curr. Biol.* **11**, 503–507.
- Saint-Jeannet, J.-P., He, X., Varmus, H. E. and Dawid, I. B. (1997). Regulation of dorsal fate in the neuraxis by *Wnt-1* and *Wnt-3a*. *Proc. Natl. Acad. Sci. USA* **94**, 13713–13718.
- Sakai, D. and Trainor, P. A. (2009). Treacher Collins syndrome: unmasking the role of *Tcof1/treacle*. *Int. J. Biochem. Cell Biol.* **41**, 1229–1232.
- Sasai, N., Mizuseki, K. and Sasai, Y. (2001). Requirement of *FoxD3*-class signaling for neural crest determination in *xenopus*. *Development* **128**, 2525–2536.
- Sauka-Spengler, T., Meulemans, D., Jones, M. and Bronner-Fraser, M. (2007). Ancient evolutionary origin of the neural crest gene regulatory network. *Dev. Cell* **13**, 405–420.
- Schorle, H., Meier, P., Buchert, M., Jaenisch, R. and Mitchell, P. J. (1996). Transcription factor *AP-2* essential for cranial closure and craniofacial development. *Nature* **381**, 235–238.
- Shi, J., Severson, C., Yang, J., Wedlich, D. and Klymkowsky, M. W. (2011). *Snail2* controls mesodermal BMP/*Wnt* induction of neural crest. *Development* **138**, 3135–3145.
- Simões-Costa, M. and Bronner, M. E. (2013). Insights into neural crest development and evolution from genomic analysis. *Genome Res.* **23**, 1069–1080.
- Simões-Costa, M. and Bronner, M. E. (2015). Establishing neural crest identity: a gene regulatory recipe. *Development* **142**, 242–257.
- Sock, E., Schmidt, K., Hermanns-Borgmeyer, I., Bösl, M. R. and Wegner, M. (2001). Idiopathic weight reduction in mice deficient in the high-mobility-group transcription factor *Sox8*. *Mol. Cell. Biol.* **21**, 6951–6959.
- Spokony, R. F., Aoki, Y., Saint-Germain, N., Magner-Fink, E. and Saint-Jeannet, J. P. (2002). The transcription factor *Sox9* is required for cranial neural crest development. *Development* **129**, 421–432.
- Steventon, B., Carmona-Fontaine, C. and Mayor, R. (2005). Genetic network during neural crest induction: from cell specification to cell survival. *Semin. Cell Dev. Biol.* **16**, 647–654.
- Steventon, B., Araya, C., Linker, C., Kuriyama, S. and Mayor, R. (2009). Differential requirements of BMP and *Wnt* signalling during gastrulation and neurulation define two steps in neural crest induction. *Development* **136**, 771–779.

- Stewart, R. A., Arduini, B. L., Berghmans, S., George, R. E., Kanki, J. P., Henion, P. D. and Look, A. T.** (2006). Zebrafish *foxd3* is selectively required for neural crest specification, migration and survival. *Dev. Biol.* **292**, 174-188.
- Sun, X., Zhang, R., Lin, X. and Xu, X.** (2008). Wnt3a regulates the development of cardiac neural crest cells by modulating expression of cysteine-rich intestinal protein 2 in rhombomere 6. *Circ. Res.* **102**, 831-839.
- Takada, S., Stark, K. L., Shea, M. J., Vassileva, G., McMahon, J. A., McMahon, A. P.** (1994). Wnt-3a regulates somite and tailbud formation in the mouse embryo. *Genes Dev.* **8**, 174-189.
- Taneyhill, L. A., Coles, E. G. and Bronner-Fraser, M.** (2007). Snail2 directly represses cadherin6B during epithelial-to-mesenchymal transitions of the neural crest. *Development* **134**, 1481-1490.
- Teng, L., Mundell, N. A., Frist, A. Y., Wang, Q. and Labosky, P. A.** (2008). Requirement for *Foxd3* in the maintenance of neural crest progenitors. *Development* **135**, 1615-1624.
- Theveneau, E., Marchant, L., Kuriyama, S., Gull, M., Moepps, B., Parsons, M. and Mayor, R.** (2010). Collective chemotaxis requires contact-dependent cell polarity. *Dev. Cell* **19**, 39-53.
- Trainor, P. A.** (2005). Specification of neural crest cell formation and migration in mouse embryos. *Semin. Cell Dev. Biol.* **16**, 683-693.
- Trainor, P. A. and Andrews, B. T.** (2013). Facial dysostoses: etiology, pathogenesis and management. *Am. J. Med. Genet. C Semin. Med. Genet.* **163**, 283-294.
- Tribulo, C., Aybar, M. J., Nguyen, V. H., Mullins, M. C. and Mayor, R.** (2003). Regulation of *Msx* genes by a *Bmp* gradient is essential for neural crest specification. *Development* **130**, 6441-6452.
- Vadasz, S., Marquez, J., Tulloch, M., Shylo, N. A. and García-Castro, M. I.** (2013). Pax7 is regulated by cMyb during early neural crest development through a novel enhancer. *Development* **140**, 3691-3702.
- van Amerongen, R. and Berns, A.** (2006). Knockout mouse models to study Wnt signal transduction. *Trends Genet.* **22**, 678-689.
- Villanueva, S., Glavic, A., Ruiz, P. and Mayor, R.** (2002). Posteriorization by FGF, Wnt, and retinoic acid is required for neural crest induction. *Dev. Biol.* **241**, 289-301.
- Yan, Y.-L., Willoughby, J., Liu, D., Crump, J. G., Wilson, C., Miller, C. T., Singer, A., Kimmel, C., Westerfield, M. and Postlethwait, J. H.** (2005). A pair of Sox: distinct and overlapping functions of zebrafish *sox9* co-orthologs in craniofacial and pectoral fin development. *Development* **132**, 1069-1083.
- Yoshikawa, Y., Fujimori, T., McMahon, A. P. and Takada, S.** (1997). Evidence that absence of Wnt-3a signaling promotes neuralization instead of paraxial mesoderm development in the mouse. *Dev. Biol.* **183**, 234-242.
- Zhang, J., Hagopian-Donaldson, S., Serbedzija, G., Elsemore, J., Plehn-Dujowich, D., McMahon, A. P., Flavell, R. A. and Williams, T.** (1996). Neural tube, skeletal and body wall defects in mice lacking transcription factor AP-2. *Nature* **381**, 238-241.